



RICE



BRIEFS

April-May 2008

In this issue...

- **Algae and phosphate**
- **Early field drainage and nitrogen losses**
- **Straw and nitrogen**

Planting season has started in California. As of April 21st, 2% of California's rice acreage had been planted but none emerged yet according to the National Agricultural Statistics Service. The unusual low temperatures registered during the last part of April may cause some delay in the emergence and growth of early plantings. This opens the door for disease and pest problems. There's not much that can be done about weather conditions, so fields should be monitored to make sure seeds are emerging normally and ensure a good stand.

Another common problem during early stages of rice growth is algae. Included in this issue of Rice Briefs is an article on algal development and its relationship with phosphorous fertilization. The information presented may not help prevent algal problems during the current season; however, it may be used to improve phosphorous fertilization practices in the near future.

Also included in this newsletter is an explanation of how early field drainage can cause nitrogen losses. Early drainage is becoming a common practice when using contact herbicides for weed control. If nitrogen losses are understood, it may be possible to mitigate these losses. Research will be conducted this season to develop fertilization strategies that could reduce nitrogen losses.

Phosphate and Rice-Field Algae

David Spencer, Ecologist, USDA-ARS Exotic and Invasive Weeds Research Unit

One approach to managing excessive algal growth is to alter the environmental conditions that support this growth. One strategy involves reducing abundance of what is often the limiting nutrient for algae in freshwater systems, phosphate. With the help of cooperators, we collected 63 water samples from rice fields and analyzed the samples for phosphate.

The results showed that phosphate levels ranged from 0.002 to 1.036 parts per million and that just over half (54%) of the water samples had phosphate levels that would limit the growth of the so-called black algae, *Nostoc spongiaeforme*. On May 9, 2007 we collected water samples from a rice field within 48 hours of flooding, but before water had been transferred out of the field. This field had phosphorus applied to the surface, but it was not incorporated into the soil. Phosphate levels in the water increased (up to five times that of the incoming water) as the water moved across a field, from south to north (Figure 1).

These results demonstrate that applying inorganic phosphorus to the field's surface without incorporating it into the soil results in much higher phosphate levels in the water. This inorganic phosphorus would directly support algal growth.

The combined results of an algal growth experiment (that showed the levels of phosphate that limited growth of *Nostoc spongiaeforme*), the measurements of phosphate levels in initial rice field water samples, and the data showing that phosphate levels in rice field water increased in a field where the phosphorus fertilizer was surface applied but not incorporated, provide strong evidence to support the notion that incorporating phosphorus fertilizer would lead to reduced phosphate levels in rice field water and subsequent reduced growth of algae, especially the so-called "black algae," *Nostoc spongiaeforme*.



Figure 1. Phosphate (mg/L) in rice field water, 48 hours after water started entering the field.

Understanding Nitrogen Losses Due to Early Field Drainage

Luis Espino, UCCE Rice Farm Advisor, and Bruce Linquist, Project Scientist, Plant Sciences Department, UC Davis

Due to recent challenges in weed control, water management during early rice growth stages has suffered significant changes. New herbicides are being used, and some of these are contact herbicides that do not translocate inside the plant killing only the part of the plant covered by the spray. To achieve good weed control with contact herbicides the field must be fully drained to expose weeds to the herbicide.

Early draining and reflooding of rice fields affect the way nitrogen (N) behaves in the soil. N management guidelines for California were developed for fields that were flooded before planting and remained flooded until harvest. Draining and reflooding fields early in the season can cause severe losses of N. To better understand how this happens, it is necessary to understand how N behaves in flooded soils.


N in flooded soils

N can be taken up by plants in two forms: NH_4^+ (ammonium ions) and NO_3^- (nitrate ions). Any fertilizer applied to the soil needs to change into any of these two N forms to be absorbed by the rice plant. Under anaerobic conditions (lack of oxygen), as in flooded rice fields, NH_4^+ is stable and accumulates, while NO_3^- is very unstable and is lost to the atmosphere. Because of this, under flooded conditions rice mostly uses NH_4^+ .

Flood-Drain-Reflood

In California, most fields are fertilized with N in the form of aqua ammonia injected into the soil before planting. Under flooded conditions, aqua produces NH_4^+ , which is taken up by the plant. When the field is drained, conditions change from anaerobic to aerobic (presence of oxygen). Under aerobic conditions, soil microbes use the NH_4^+ and turn it into NO_3^- . This process is called **nitrification**. Nitrification does not change the amount of N in the soil, just its form. Later, when the field is reflooded, the soil microorganisms can't use oxygen anymore, and instead use NO_3^- , transforming it into NO_2 (nitrous oxide) and other N gases that can be lost to the atmosphere. This process is called **denitrification**.

The whole process is summarized in the following diagram:



Thus, draining a field does not cause loss of N, but just accumulation of NO_3^- . N is only lost after the field is reflooded.

Reducing N loss due to reflooding

Research conducted during 2006 and 2007 at the Rice Experiment Station (Biggs) and in commercial rice fields with different soil types and straw management practices has yielded very interesting results. Experiments showed that during the drain period N in the form of NO_3^- accumulates at an average rate of 1.8 lbs/acre/day. As explained above, after reflooding this N is susceptible to losses due to denitrification. It only took 7 days for the accumulated NO_3^- to disappear from the soil. N uptake and yield reductions were observed in some of the fields that were flooded-drained-reflooded, most likely due to N losses or/and drought stress caused during the drain.

These results show that by minimizing the length of the drain period as much as possible significant N losses can be avoided. Research will continue in the coming years to develop improved strategies of N application in fields where early drains are used.

Straw Incorporation and Nitrogen Management*

The reduction of rice straw burning brought many changes in the way straw is managed in California rice fields. Research has been conducted to determine how changes in straw management impact other areas of rice production, such as soil fertility. In many fields rice straw is being incorporated after grain harvest while in others straw is being removed and used out of the field. Research has shown that the incorporation of rice straw can have beneficial effects in soil fertility. A study conducted in the Sacramento valley concluded that after 5 years of straw incorporation, N application rates can be reduced 25 lbs/acre.

Rice straw can have significant amounts of macro and micro nutrients:

Nutrient content of rice straw in lbs/acre (assumes 10,000 lbs of straw/acre)

| N | P | K | S | Ca | Mg | Na | Cl | B | Zn | Cu | Mo |
|----|----|----|----|----|----|----|----|------|------|------|------|
| 68 | 14 | 77 | 10 | 28 | 18 | 78 | 8 | 0.15 | 0.41 | 0.31 | 0.02 |

When incorporating rice straw, many of the nutrients extracted by the rice plant are being returned to the soil and, after the straw is decomposed, will be available for plant uptake. Straw incorporation increases microbial biomass N, that is, N being used by microbes in partially broken-down straw residues. It is this portion of N that will be readily available for crop uptake.

After a few years of straw incorporation, a fraction of the N from synthetic fertilizers (fertilizer N) applied to the field is “captured” by soil microorganisms. These microorganisms compete with the plant to use the fertilizer N. At the same time, N is being released from the microbial biomass formed by the incorporation of rice straw, and this N replaces the fertilizer N in the soil solution and is taken up by the rice plant. Research has shown that when straw is incorporated more fertilizer N will be available after one season of being applied than when straw is removed.

Straw incorporation also can affect weed density. When compared to burned fields, fields where straw was incorporated had higher density of grassy weeds, especially watergrass. However, when fields were winter flooded, watergrass density was reduced. It is thought that this reduction may be due to the foraging activity of waterfowl.

*Adapted from Bird, J. A., A. J. Eagle, W. R. Horwath, M. W. Hair, E. E. Zilbert and C. van Kessel. 2002. Long-term studies find benefits, challenges in alternative rice straw management. *California Agriculture* 56: 69-75.

To receive this newsletter electronically or for suggestions/questions send an email to: laespino@ucdavis.edu, or go to <http://cecolusa.ucdavis.edu/rice/>



This newsletter is produced by:

Luis Espino

Farm Advisor

P.O. Box 180
100 Sunrise Blvd., Suite E

Colusa, CA 95932

(530) 458-0570

(530) 635-6234

Email: laespino@ucdavis.edu

<http://cecolusa.ucdavis.edu/rice/>



To simplify information, tradenames of products have been used. No endorsements of named products is intended, nor is criticism of similar products which are not mentioned.

The University of California prohibits discrimination or harassment of any person on the basis of race, color, national origin, religion, sex, gender identity, pregnancy (including childbirth, and medical conditions related to pregnancy or childbirth), physical or mental disability, medical condition (cancer-related or genetic characteristics), ancestry, marital status, age, sexual orientation, citizenship, or status as a covered veteran (covered veterans are special disabled veterans, recently separated veterans, Vietnam era veterans, or any other veterans who served on active duty during a war or in a campaign or expedition for which a campaign badge has been authorized) in any of its programs or activities. University policy is intended to be consistent with the provisions of applicable State and Federal laws. Inquiries regarding the University's nondiscrimination policies may be directed to the Affirmative Action/Staff Personnel Services Director, University of California, Agriculture and Natural Resources, 300 Lakeside Drive, 6th Floor, Oakland, CA 94612-3550, (510) 987-0096.

Co-operative Extension Work in Agriculture and Home Economics U.S. Department of Agriculture, University of California, and County of Colusa cooperating.

COOPERATIVE EXTENSION

University of California

P.O. Box 180

Colusa, CA 95932

Non-Profit Org.

U.S. POSTAGE

P A I D

Colusa, CA

95932

Permit No. 3

Rice Briefs